



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB09850 - Goat Anti-CRISP2 (aa77-89) Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** CRISP-2, cysteine-rich secretory protein 2, GAPDL5, glyceraldehyde-3-phosphate dehydrogenase-like 5, MGC111136, OTTHUMP00000016589, testis specific protein 1 (probe H4-1 p3-1), TPX1, TSP1, CRISP2

**Official Symbol:** CRISP2

**Accession Number(s):** NP\_003287.1

**Human GeneID(s):** [7180](#)

**Important Comments:** Reported variants represent identical protein: NP\_003287.1, NP\_001135879.1, NP\_001135889.1, NP\_001135907.1, NP\_001135880.1.

### Immunogen

Peptide with sequence KCTLQHSDPEDRK, from the internal region of the protein sequence according to NP\_003287.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested


**Peptide ELISA:** antibody detection limit dilution 1:128000.

**Western blot:** In transfected HEK293 transiently expressing Human CRISP2 (MYC and DYKDDDDK tagged), a band of approx. 35kDa was observed. No bands were observed in mock-transfected HEK293 and the same band was observed using anti-myc tag antibody. Recommended concentration, 0.05-0.1µg/ml.

### Species Reactivity

**Tested:**

**Expected from sequence similarity:** Human, Pig, Dog, Cow



HEK293 lysate (10ug protein in RIPA buffer) overexpressing Human CRISP2 (RC205312) with C-terminal MYC tag probed with EB09850 (0.11ug/ml) in Lane A and probed with anti-MYC Tag (1/1000) in lane C. Mock-transfected HEK293 probed with EB09850 (1mg/ml) in Lane B. Primary incubations were for 1 hour. Detected by chemiluminescence.