



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB08182 - Goat Anti-CRP2 / CSRP2 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** CRP2, CSRP2, cysteine and glycine-rich protein 2, LMO5, SmLIM, LIM domain only 5, smooth muscle

**Official Symbol:** CSRP2

**Accession Number(s):** NP\_001312.1

**Human GeneID(s):** [1466](#)

**Non-Human GeneID(s):** 13008 (mouse)

### Immunogen

Peptide with sequence C-ERLGKIPESVQP, from the internal region of the protein sequence according to NP\_001312.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:64000.


**Western blot:** Approx 26kDa band observed in Human, Mouse and Rat Kidney lysates (calculated MW of 21.0kDa according to NP\_001312.1). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Huber et al, J Biol Chem. 2000 Feb 25;275(8):5504-11; PMID: 10681529). In transfected HEK293 transiently expressing CRP2 a band of approx. 26kDa is observed. This band is not observed in the non-transfected HEK293. Recommended concentration: 0.1-0.3µg/ml. Primary incubation was 1 hour.

**IHC:** In paraffin embedded Human Lung shows staining of the endothelial cells of a blood vessel. Recommended concentration: 3-6µg/ml.


### Species Reactivity

**Tested:** Human, Mouse, Rat


**Expected from sequence similarity:** Human, Mouse, Rat, Horse



EB08182 (0.2 $\mu$ g/ml) staining of Human (A), Mouse (B) and Rat (C) Kidney lysates (35 $\mu$ g protein in RIPA buffer).  
Detected by chemiluminescence



HEK293 overexpressing CRP2 (RC201565) and probed with EB08182 (mock transfection in first lane), tested by  
Origene.



EB08182 (3.8µg/ml) staining of paraffin embedded Human Lung. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.