



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB08525 - Goat Anti-CTDSPL Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** CTDSPL, CTD (carboxy-terminal domain, RNA polymerase II, polypeptide A) small phosphatase-like, C3orf8, HYA22, PSR1, SCP3, small CTD phosphatase 3

**Official Symbol:** CTDSPL

**Accession Number(s):** NP\_001008393.1; NP\_005799.2

**Human GeneID(s):** [10217](#)

**Important Comments:** This antibody is expected to recognise both isoforms (NP\_001008393.1; NP\_005799.2).

### Immunogen

Peptide with sequence QCNVSLKKQRSRS, from the internal region of the protein sequence according to NP\_001008393.1; NP\_005799.2.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Preliminary testing showed a band at approx 28kDa in nuclear lysate of cell lines A431 and HeLa, after 1-2µg/ml antibody staining (calculated Mwt of 29.9kDa according to NP\_005799.2). Primary incubation 1 hour at room temperature.


**Immunofluorescence:** Strong expression of the protein seen in the nuclei and vesicles of A431 cells. Recommended concentration: 10µg/ml.

**Flow Cytometry:** Flow cytometric analysis of A431 cells. Recommended concentration: 10ug/ml.


### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Dog



EB08525 Immunofluorescence analysis of paraformaldehyde fixed A431 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear and vesicle staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB08525 Flow cytometric analysis of paraformaldehyde fixed A431 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.