



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB08878 - Goat Anti-CTLA + CTLB Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: CLTA, CTLB, clathrin, light chain (Lca), LCA, clathrin, light polypeptide (Lca), clathrin, light polypeptide A, clathrin, light chain (Lcb), LCB

Official Symbol: CLTA

Accession Number(s): NP_001824.1; NP_009027.1; NP_001070145.1; NP_001825.1; NP_009028.1

Human GeneID(s): [1211](#) , [1212](#)

Non-Human GeneID(s): 12757 (mouse), 83800 (rat)

Important Comments: This antibody is expected to recognize all reported isoforms of light chains a and b.

Immunogen

Peptide with sequence CDFNPKSSKQAKD, from the internal region of the protein sequence according to NP_001824.1; NP_009027.1; NP_001070145.1; NP_001825.1; NP_009028.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.


Western blot: Approx. 35kDa and 22-23kDa bands observed in Human Brain (Frontal Cortex) lysates (calculated MW of 23.2kDa according to NP_001825.1). The observed molecular weight of ~35kDa corresponds to earlier findings in literature with different antibodies (Okamoto et al, Am J Physiol Cell Physiol. 2000 Sep;279(3):C833-51.; PMID: 10942733). Recommended concentration: 0.1-0.3µg/ml. Primary incubation was 1 hour.

IHC: Paraffin embedded Human Spleen. Recommended concentration: 2.5µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow



EB08878 (0.1µg/ml) staining of Human Frontal Cortex lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB08878 (2.5µg/ml) staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.