

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB12525 - Goat Anti-CXCR1 (aa26-38) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: CXCR1, chemokine (C-X-C motif) receptor 1, C-C, C-C-CKR-1, CD128, CD181, CDw128a, CKR-1, CMKAR1, IL8R1, IL8RA, IL8RBA, C-X-C chemokine receptor type 1, CXC-R1, CXCR-1, IL-8 receptor type 1, IL-8R A, high affinity interleukin-8 receptor A, interleukin 8 re

Official Symbol: CXCR1

Accession Number(s): NP_000625.1

Human GeneID(s): [3577](#)

Immunogen

Peptide with sequence DYSPCMLETETLN, from the internal region (near N terminus) of the protein sequence according to NP_000625.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 39kDa band observed in Human Breast cancer lysates (calculated MW of 39.8kDa according to NP_000625.1). Recommended concentration: 0.3-1µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB12525 (0.3 μ g/ml) staining of Human Breast cancer lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.