



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB10355 - Goat Anti-CYP2E1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: CPE1, CYP2E, cytochrome P450 2E1, cytochrome P450, family 2, subfamily E, polypeptide 1, cytochrome P450, subfamily IIE (ethanol-inducible), polypeptide 1, flavoprotein-linked monooxygenase, microsomal monooxygenase, P450C2E, P450-J, xenobiotic mo, CYP2E1

Official Symbol: CYP2E1

Accession Number(s): NP_000764.1

Human GeneID(s): [1571](#)

Immunogen

Peptide with sequence C-RKVIKNVAEVK, from the internal region of the protein sequence according to NP_000764.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 50kDa band observed in Human Liver lysates (calculated MW of 56.8kDa according to NP_000764.1). Recommended concentration: 0.05-0.2µg/ml.

IHC: In paraffin embedded Human Liver shows ER-like staining in hepatocytes. Recommended concentration, 3-5µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB10355 (0.05 μ g/ml) staining of Human Liver lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB10355 (3.8 μ g/ml) staining of paraffin embedded Human Liver. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.