

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB06077 - Goat Anti-DAB2 / DOC2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: DAB2, DOC2, disabled homolog 2, mitogen-responsive phosphoprotein (Drosophila), DOC-2, mitogen-responsive phosphoprotein, disabled (Drosophila) homolog 2 (mitogen-responsive phosphoprotein), disabled homolog 2, mitogen-responsive phosphoprotein, FLJ26626

Official Symbol: DAB2

Accession Number(s): NP_001334.2

Human GenelD(s): 1601

Immunogen

Peptide with sequence C-SEMYRDPFGNPFA, from the C Terminus of the protein sequence according to NP $_$ 001334.2.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 75kDa band observed in human kidney lysates (calculated MW of 82.6kDa according to NP_001334.1). Recommended concentration: 0.01-0.03µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human

250kDa 150kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa 15kDa

EB06077 staining (0.003μg/ml) of human kidney lysate (RIPA buffer, 35μg total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.