

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB12715 - Goat Anti-DC-SIGN / CD209 Antibody

Size: 100µg specific antibody in 200µl



# **Target Protein**

**Principal Names:** CD209, CD209 molecule, CDSIGN, CLEC4L, DC-SIGN, DC-SIGN1, C-type lectin domain family 4 member L, C-type lectin domain family 4, member L, CD209 antigen, HIV gpl20-binding protein, dendritic cell-specific ICAM-3-grabbing non-integrin 1, dendritic cell-specific intracellular adhesion molecules (ICAM)-3 grabbing non-integrin

Official Symbol: CD209

**Accession Number(s):** NP\_066978.1; NP\_001138368.1; NP\_001138369.1; NP\_001138365.1; NP\_001138366.1; NP\_001138367.1; NP\_001138371.1

Human GenelD(s): 30835

**Important Comments:** This antibody is expected to recognize all reported isoforms.

#### **Immunogen**

Peptide with sequence C-NRGEPNNVGEED, from the internal region of the protein sequence according to NP\_066978.1; NP\_001138368.1; NP\_001138369.1; NP\_001138365.1; NP\_001138366.1; NP\_001138367.1; NP\_001138371.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

#### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:128000.

**Western blot:** Approx. 50kDa band observed in lysates of cell lines K562 and THP-1, and in preliminary testing of Human Lung and Testes lysate (calculated MW of 45.8kDa according to NP\_066978.1). This molecular weight is observed by other commercial sources, and was successfully blocked by incubation with the immunizing peptide. Recommended concentration: 1-3µg/ml. Primary incubation 1 hour at room temperature.

**Positive Control:** A batch specific positive control lysate is available for this product. Please contact Sales@everestbiotech.com for availability.

IHC: Paraffin embedded Human Placenta. Recommended concentration: 6µg/ml.


**Immunofluorescence:** Strong expression of the protein seen in the plasma membranes and cytoplasm of THP-1 cells. Recommended concentration: 10µg/ml.

**Flow Cytometry:** Flow cytometric analysis of THP-1 cells. Recommended concentration: 10ug/ml.


### **Species Reactivity**

Tested: Human


Expected from sequence similarity: Human




EB12715 (1 $\mu$ g/ml) staining of K562 (A) and (2 $\mu$ g/ml) THP-1 (B) cell lysate (35 $\mu$ g protein in RIPA buffer). Detected by chemiluminescence.



EB12715 (6µg/ml) staining of paraffin embedded Human Placenta. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.




EB12715 Negative Control showing staining of paraffin embedded Human Placenta, with no primary antibody.



EB12715 Immunofluorescence analysis of paraformaldehyde fixed THP-1 immobilized on Shi-fix<sup>™</sup> plus cover-slips. Primary incubation 1hr (1:50 dilution) followed by Alexa Fluor® 488 secondary antibody (1:2000 dilution), showing membrane and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control:

Anti-Goat IgG followed by Alexa Fluor® 488 secondary antibody.



EB12715 Flow cytometric analysis of paraformaldehyde fixed THP1 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control:

Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.