

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB10446 - Goat Anti-DCTN3 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: DCTN22, DCTN-22, dynactin 3, dynactin 3 (p22), dynactin 3, isoform

1, dynactin light chain, MGC111190, DCTN3

Official Symbol: DCTN3

Accession Number(s): NP_009165.1; NP_077324.1

Human GeneID(s): 11258

Non-Human GenelD(s): 53598 (mouse), 362504 (rat)

Important Comments: This antibody is expected to recognize both reported isoforms

(NP_009165.1; NP_077324.1).

Immunogen

Peptide with sequence C-QVALGNISSKRE, from the internal region of the protein sequence according to NP_009165.1; NP_077324.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 20kDa band observed in Human Liver lysates, and approx. 18kDa in Rat Kidney lysates and in preliminary testing of Mouse Kidney lysate (calculated MW of 19.5kDa according to Human NP_077324.1 and 21.1kDa according to Rat

NP_001102129.1. Recommended concentration: 0.1-0.3µg/ml. Primary incubation 1 hour at room temperature.

IHC: Paraffin embedded Human Kidney. Recommended concentration: 6-8µg/ml.

Flow Cytometry: Flow cytometric analysis of K562 cells. Recommended concentration: 10ug/ml.


Species Reactivity

Tested: Human, Rat


Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa


EB10446 (0.1μg/ml) staining of Human Liver lysate (35μg protein in RIPA buffer). Detected by chemiluminescenceWS10446-4




 $EB10446\ (0.3\mu\text{g/ml})\ staining\ of\ Rat\ Kidney\ lysate\ (35\mu\text{g}\ protein\ in\ RIPA\ buffer).\ Detected\ by\ chemiluminescence$



EB10446 (8µg/ml) staining of paraffin embedded Human Kidney. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



EB10446 Negative Control showing staining of paraffin embedded Human Kidney, with no primary antibody.



EB10446 Flow cytometric analysis of paraformaldehyde fixed K562 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control:

Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.