

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB07126 - Goat Anti-DEAD-box protein 6 Antibody

Size: 100µg specific antibody in 200µl

Target Protein

Principal Names: DEAD-box protien 6, DEAD (Asp-Glu-Ala-Asp) box polypeptide 6, DDX6, HLR2, P54, RCK, DEAD box-6, DEAD/H (Asp-Glu-Ala-Asp/His) box polypeptide 6

(RNA helicase, 54kD)

Official Symbol: DDX6

Accession Number(s): NP_004388.1

Human GeneID(s): 1656

Immunogen

Peptide with sequence C-AEYHSEPVEDEKP, from the C Terminus of the protein sequence according to NP_004388.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 50kDa band observed in lysates of cell lines Daudi, Jurkat and HepG2 (calculated MW of 53.2kDa according to NP_004388.1). Recommended

concentration: 0.3-1µg/ml. Primary incubation was 1 hour.


IHC: Paraffin embedded Human Brain (Cortex) and Thymus. Recommended

concentration: 5µg/ml.


Species Reactivity

Tested: Human


Expected from sequence similarity: Human, Mouse, Rat, Dog



EB07126 (0.3μg/ml) staining of Daudi (A), Jurkat (B) and HepG2 (C) lysates (35μg protein in RIPA buffer). Detected by chemiluminescence.



EB07126 (5µg/ml) staining of paraffin embedded Human Cortex. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB07126 ($5\mu g/ml$) staining of paraffin embedded Human Thymus. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.