

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB09742 - Goat Anti-DEFA1/3 precursor Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** DEFA1, defensin, alpha 1, myeloid-related sequence, HGNC:2761, DEF1, HNP-1, HP-1, MRS, defensin, alpha 1, myeloid-related sequence, DEFA3,

defensin, alpha 3, neutrophil-specific, neutrophil peptide 3

Official Symbol: DEFA1

Accession Number(s): NP\_004075.1; NP\_005208.1

Human GenelD(s): 1667, 1668

Important Comments: This antibody is expected to equally recognize DEFA1 precursor

and DEFA3 precursor.

## Immunogen

Peptide with sequence C-DESLAPKHPGSRKN, from the internal region of the protein sequence according to NP\_004075.1; NP\_005208.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:128000.

**Western blot:** Approx 10kDa band observed in Human Spleen lysates (calculated MW of 10.2kDa according to NP\_004075.1). An additional band of unknown identity was also consistently observed at 20kDa, and was successfully blocked by incubation with the immunizing peptide. Recommended concentration: 1-3µg/ml. Primary incubation 1 hour at room temperature.

IHC: Paraffin embedded Human Thymus. Recommended concentration: 3-6µg/ml.


# **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB09742 (2μg/ml) staining of Human Spleen lysate (35μg protein in RIPA buffer). Detected by chemiluminescence.



EB09742 (3.8 $\mu$ g/ml) staining of paraffin embedded Human Thymus. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.