

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB10758 - Goat Anti-Delta-Sarcoglycan (aa245-257) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: SGCD, sarcoglycan, delta (35kDa dystrophin-associated glycoprotein), 35DAG, CMD1L, DAGD, MGC22567, SG-delta, SGCDP, SGD, 35kD

dystrophin-associated glycoprotein, delta-SG, delta-sarcoglycan, dystrophin associated

Official Symbol: SGCD

Accession Number(s): NP_000328.2; NP_758447.1; NP_001121681.1

Human GeneID(s): 6444

Non-Human GenelD(s): 24052 (mouse), 497892 (rat)

Important Comments: This antibody is expected to recognize isoform 1 (NP_000328.2)

and isoform 3 (NP_001121681.1).

Immunogen

Peptide with sequence C-RLPHGSYTPTGTR, from the internal region of the protein sequence according to NP_000328.2; NP_758447.1; NP_001121681.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 37kDa band observed in Human and Rat Heart lysates (calculated MW of 32.2kDa according to NP_000328.2). The observed molecular weight corresponds to the glycosylated form. Recommended concentration: 0.1-0.3µg/ml.

IHC: In paraffin embedded Human Skeletal Muscle shows sarcolemmar staining of the fibres. Recommended concentration, 3-5µg/ml.

Species Reactivity

Tested: Human, Rat

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow

250kDa 150kDa 100kDa 75kDa 50kDa


37kDa

25kDa

20kDa

15kDa

EB10758 (0.1 μ g/ml) staining of Human Heart lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB10758 (3.8μg/ml) staining of paraffin embedded Human Skeletal Muscle. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.