

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB09106 - Goat Anti-Desmin Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: DES, desmin, CMD1I, CSM1, CSM2, FLJ12025, FLJ39719, FLJ41013,

FLJ41793, intermediate filament protein

Official Symbol: DES

Accession Number(s): NP_001918.3

Human GeneID(s): 1674

Non-Human GenelD(s): 13346 (mouse), 64362 (rat)

Immunogen

Peptide with sequence C-RDGEVVSEATQQQHE, from the C Terminus of the protein sequence according to NP_001918.3.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx. 55-60kDa band observed in Human and Mouse Skeletal Muscle lysates and in Human Heart lysates (calculated MW of 53.5kDa according to Human NP_001918.3, and Mouse NP_034173.1). Preliminary testing showed a band at approx 60kDa in Rat and Pig Skeletal muscle lysate. Recommended concentration:


0.01-0.3µg/ml. Primary incubation 1 hour at room temperature.

IHC: Paraffin embedded Human Heart. Recommended concentration: 4-6µg/ml.

Species Reactivity

Tested: Human, Mouse


Expected from sequence similarity: Human, Mouse, Dog, Cow




EB09106 ($0.01\mu g/ml$) staining of Human Skeletal Muscle (A) and (0.3ug/ml) Heart (B) lysate ($35\mu g$ protein in RIPA buffer). Detected by chemiluminescence.



EB09106 (0.1μg/ml) staining of Mouse Skeletal Muscle lysate (35μg protein in RIPA buffer). Detected by chemiluminescence.



EB09106 ($6\mu g/ml$) staining of paraffin embedded Human Heart. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



EB09106 Negative Control showing staining of paraffin embedded Human Heart, with no primary antibody.