

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB07575 - Goat Anti-DGAT1 Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** DGAT1, diacylglycerol O-acyltransferase homolog 1 (mouse), ARGP1, DGAT, ACAT related gene product 1, acyl coenzyme A:cholesterol acyltransferase related

gene 1, diacylglycerol O-acyltransferase 1, diglyceride acyltransferase

Official Symbol: DGAT1

Accession Number(s): NP\_036211.2

Human GeneID(s): 8694

Non-Human GenelD(s): 13350 (mouse), 84497 (rat)

#### **Immunogen**

Peptide with sequence C-QNSMKPFKDMDYS, from the internal region of the protein sequence according to NP\_036211.2.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:64000.

**Western blot:** Approx 55kDa band observed in Mouse Liver lysates (calculated MW of 56.8kDa according to NP\_034176.1). Recommended concentration: 0.01-0.1.0μg/ml. Primary incubation 1 hour at room temperature.

IHC: Paraffin embedded Human Colon. Recommended concentration: 3.75µg/ml.

**Immunofluorescence:** Strong expression of the protein seen in the endoplasmic reticulum and cytoplasm of A431 cells. Recommended concentration: 10µg/ml.

**Flow Cytometry:** Flow cytometric analysis of A431 cells. Recommended concentration: 10ug/ml.

#### **Species Reactivity**

Tested: Human, Mouse


Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa


EB07575 (0.03μg/ml) staining of Mouse Liver lysate (35μg protein in RIPA buffer). Detected by chemiluminescence.



EB07575 Immunofluorescence analysis of paraformaldehyde fixed A431 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing endoplasmic reticulum and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control:



EB07575 (3.75 $\mu$ g/ml) staining of paraffin embedded Human Colon. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB07575 Flow cytometric analysis of paraformaldehyde fixed A431 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control:

Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.