

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB07661 - Goat Anti-DGAT2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: DGAT2, diacylglycerol O-acyltransferase homolog 2 (mouse),

DKFZp686A15125, HMFN1045, GS1999full, diacylglycerol O-acyltransferase homolog 2,

diacylglycerol O-acyltransferase-like protein 2 isoform 1

Official Symbol: DGAT2

Accession Number(s): NP_115953.2; NP_001240820.1

Human GeneID(s): 84649

Non-Human GenelD(s): 67800 (mouse), 252900 (rat)

Important Comments: This antibody is not expected to cross-react with other DGAT

members.

Immunogen

Peptide with sequence C-KLEHPTQQDIDLYH, from the internal region (near the C Terminus) of the protein sequence according to NP_115953.2; NP_001240820.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

IHC: In paraffin embedded Human Liver shows granular cytoplasm staining of

hepatocytes. Recommended concentration: 3µg/ml.


Flow Cytometry: Flow cytometric analysis of HeLa cells. Recommended concentration:

10ug/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog



EB07661 (3µg/ml) staining of paraffin embedded Human Liver. Microwaved antigen retrieval with Tris/EDTA buffer pH9, HRP-staining.



EB07661 Flow cytometric analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control:

Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.