



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05518 - Goat Anti-Dicarbonyl Reductase Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: DCXR, dicarbonyl/L-xylulose reductase, DCR, HCR11, KIDCR, P34H, carbonyl reductase, carbonyl reductase II, kidney dicarbonyl reductase, HCR2, SDR20C1, human carbonyl reductase 2, kidney dicarbonyl reductase, short chain dehydrogenase/reductase family 20C, member 1

Official Symbol: DCXR

Accession Number(s): NP_057370.1

Human GeneID(s): [51181](#)

Immunogen

Peptide with sequence GSTLPVEGGFWAC, from the C Terminus of the protein sequence according to NP_057370.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 26kDa band observed in human kidney, liver and testes lysates (calculated MW of 25.9kDa according to NP_057370.1). Recommended for use at 0.1-0.3µg/ml.

IHC: In paraffin embedded human kidney, stains membrane / cytoplasm in epithelial cells of renal glomeruli and tubules. Recommended concentration, 1-2µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB05518 staining (0.1 μ g/ml) of human kidney lysate (RIPA buffer, 35 μ g total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.



EB05518 (1 μ g/ml) staining of paraffin embedded human kidney. Microwaved antigen retrieval with citrate buffer pH 6, HRP-staining.