

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB11466 - Goat Anti-Dishevelled1 (aa20-32) Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** Dvl1, dishevelled, dsh homolog 1 (Drosophila), RP23-242K15.15-003, Dvl, mKIAA4029, DSH homolog 1, dishevelled-1, segment polarity protein dishevelled

homolog DVL-1

Official Symbol: Dvl1

Accession Number(s): NP\_034221.3

Human GenelD(s): 1855

Non-Human GenelD(s): 13542 (mouse), 83721 (rat)

## **Immunogen**

Peptide with sequence KLPVAPERVTLAD-C, from the N Terminus of the protein sequence according to NP\_034221.3.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

#### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:128000.

IHC: Paraffin embedded Human Skeletal Muscle and Kidney. Recommended

concentration: 5µg/ml.

## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Pig, Cow

#### **Specific Reference**


Spate LD, Brown AN, Redel BK, Whitworth KM, Murphy CN, Prather RS.

Dickkopf-related protein 1 inhibits the WNT signaling pathway and improves pig oocyte


maturation.

PLoS One. 2014 Apr 16;9(4):e95114.

PMID: 24739947



EB11466 ( $5\mu g/ml$ ) staining of paraffin embedded Human Skeletal Muscle. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB11466 (5μg/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.