



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB08317 - Goat Anti-DNMT1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: DNMT1, DNA (cytosine-5-)-methyltransferase 1, CXXC9, DNMT, FLJ16293, MCMT, MGC104992, DNA methyltransferase 1

Official Symbol: DNMT1

Accession Number(s): NP_001370.1

Human GeneID(s): [1786](#)

Non-Human GeneID(s): 13433 (mouse), 84350 (rat)

Immunogen

Peptide with sequence C-RFESPPKTQPTEDN, from the internal region of the protein sequence according to NP_001370.1 .

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 170kDa band observed in lysates of cell line Jurkat (calculated MW of 183kDa according to NP_001370.1). Recommended concentration: 0.5-1.5µg/ml. Primary incubation was 1 hour. Preliminary testing was unsuccessful on Mouse for this particular batch.

IHC: In paraffin embedded Human Kidney shows nuclear staining in PCT. Recommended concentration: 3-6µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow, Pig



EB08317 (0.5µg/ml) staining of Jurkat cell lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB08317 (3.8µg/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.