



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB09820 - Goat Anti-DYDC1 (aa142-154) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: bA36D19.5, BC019250 protein, DPY30 domain containing 1, DPY30D1, FLJ43920, OTTHUMP00000019963, DYDC1

Official Symbol: DYDC1

Accession Number(s): NP_620167.1

Human GeneID(s): [143241](#)

Non-Human GeneID(s): 69496 (mouse), 688433 (rat)

Immunogen

Peptide with sequence C-TLAEISDRYGAPN, from the internal region of the protein sequence according to NP_620167.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: n transfected HEK293 transiently expressing Human DYDC1 (myc and DYKDDDDK tagged), a band of approx. 28kDa is observed. No bands are observed in mock-transfected HEK293 and the same band is observed using anti-myc tag antibody. This 28kDa band is also observed in Human, Mouse and Rat Testis lysates.


Recommended concentration: 0.1-1µg/ml.

IHC: Paraffin embedded Human Testis. Recommended concentration: 5µg/ml.


Species Reactivity

Tested: Human, Mouse, Rat

Expected from sequence similarity: Human, Mouse, Rat, Cow



Lane A - EB09820 (1 μ g/ml) staining of HEK293 overexpressing Human DYDC1 lysate (10 μ g protein in RIPA buffer) Lane B - EB09820 (0.1 μ g/ml) staining of HEK293 mock-transfected lysate (10 μ g protein in RIPA buffer). Lane C - EB00106 anti-MYC Tag (1/1000) staining HEK293 overexpressing Human DYDC1 lysate (10 μ g protein in RIPA buffer). Primary incubations were for 1 hour. Detected by chemiluminescence.



EB09820 (5 μ g/ml) staining of paraffin embedded Human Testis. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.