

### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB05367 - Goat Anti-Dysadherin Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** FXYD5, dysadherin, FXYD domain containing ion transport regulator 5, RIC, IWU1, IWU-1, dysad, HSPC113, KCT1, OIT2, PRO6241, FXYD domain-containing ion transport regulator 5, keratinocytes associated transmembrane protein 1, DYSAD,

HSPC113

Official Symbol: FXYD5

Accession Number(s): NP\_054883.3; NP\_659003.1; NP\_001158077.1

Human GeneID(s): 53827

Important Comments: NP\_054883.3, NP\_659003.1and NP\_001158077.1 are variants

that represent the same protein.

## Immunogen

Peptide with sequence GKCRQLSRLCRNHCR, from the C Terminus of the protein sequence according to NP\_054883.3; NP\_659003.1; NP\_001158077.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:8000.

**Western blot:** Approx 38kDa band observed in Human Spleen lysates (calculated MW of 19.5 kDa according to NP\_054883 and NP\_659003). The band was successfully blocked by incubation with the immunizing peptide. According to the literature this protein is O-glycosylated (PMID: 12672699), which may explain the size difference observed.

## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human



EB05367 staining (0.5μg/ml) of Human Spleen lysate (RIPA buffer, 30μg total protein per lane). Primary incubated for 1 hour. Detected by chemiluminescence.