

## **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB05995 - Goat Anti-ECT2 Antibody

Size: 100µg specific antibody in 200µl



# **Target Protein**

Principal Names: ECT2, epithelial cell transforming sequence 2 oncogene, FLJ10461,

epithelial cell transforming sequence 2 oncogene protein

Official Symbol: ECT2

**Accession Number(s):** NP\_060568.3; NP\_001245244.1; NP\_001336023.1; NP\_001336026.1; NP\_001336030.1; NP\_001336031.1; NP\_001336032.1

Human GeneID(s): 1894

Important Comments: This antibody is expected to recognise NP\_001245244.1 (iso a),

NP\_060568.3 (iso b), NP\_001336023.1 (iso c), NP\_001336026.1 (iso d), NP\_001336030.1 (iso e), NP\_001336031.1 (iso f); NP\_001336032.1 (iso g)

# **Immunogen**

Peptide with sequence AENSVLTSTTGRT-C, from the N Terminus of the protein sequence according to NP\_060568.3; NP\_001245244.1; NP\_001336023.1; NP\_001336026.1; NP\_001336030.1; NP\_001336031.1; NP\_001336032.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:128000.


**Immunofluorescence:** Strong expression of the protein seen in the nuclei and cytoplasm of A431 and U2OS cells. Recommended concentration: 10µg/ml.

**Flow Cytometry:** Flow cytometric analysis of A431 cells. Recommended concentration: 10ug/ml.


## **Species Reactivity**

Tested: Human


Expected from sequence similarity: Human



EB05995 Immunofluorescence analysis of paraformaldehyde fixed A431 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing strong nuclear and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB05995 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing strong nuclear and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB05995 Flow cytometric analysis of paraformaldehyde fixed A431 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control:

Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.