

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB11689 - Goat Anti-EEA1 (aa821-835) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: early endosome antigen 1, early endosome antigen 1, 162kD, early endosome-associated protein, EEA1, endosome-associated protein p162, MST105, MSTP105, ZFYVE2, zinc finger FYVE domain-containing protein 2

Official Symbol: EEA1

Accession Number(s): NP_003557.2

Human GeneID(s): 8411

Immunogen

Peptide with sequence C-QETKIQHEELNNRIQ, from the internal region of the protein sequence according to NP_003557.2.

Please note the <u>peptide</u> is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 200kDa band observed in lysates of cell line HEK293 (calculated MW of 162kDa according to NP_003557.2). This molecular weight is routinely observed by other sources. Recommended concentration: $0.3-1\mu g/ml$. Primary incubation 1 hour at room temperature.

Immunofluorescence: Strong expression of the protein seen in the vesicles/cytoplasm of NIH3T3 cells. Recommended concentration: 5µg/ml.


Species Reactivity

Tested: Human, Mouse

Expected from sequence similarity: Human

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB11689 (0.3μg/ml) staining of HEK293 lysate (35μg protein in RIPA buffer). Detected by chemiluminescence.



EB11689 Immunofluorescence analysis of paraformaldehyde fixed NIH3T3 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (5ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing vesicle/cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (5ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).