



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB12188 - Goat Anti-EEF1E1 / AIMP3 / p18 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** EEF1E1, eukaryotic translation elongation factor 1 epsilon 1, AIMP3, P18, ARS-interacting multifunctional protein 3, aminoacyl tRNA synthetase complex-interacting multifunctional protein 3, eukaryotic translation elongation factor 1 epsilon-1,

**Official Symbol:** EEF1E1

**Accession Number(s):** NP\_004271.1; NP\_001129122.1

**Human GeneID(s):** [9521](#)

**Important Comments:** This antibody is expected to recognize both reported isoforms (NP\_004271.1; NP\_001129122.1).

### Immunogen

Peptide with sequence C-SKGNKYSAQGERQ, from the internal region (near N Terminus) of the protein sequence according to NP\_004271.1; NP\_001129122.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:128000.


**Western blot:** Approx 19kDa band observed in lysates of cell lines HepG2 and K562 (calculated MW of 19.8kDa according to NP\_004271.1) and in Pig Testis lysates (calculated MW of 19.8kDa according to NP\_001230474.1). Recommended concentration: 0.1-0.3µg/ml. Primary incubation was 1 hour.

**IHC:** Paraffin embedded Human Pancreas. Recommended concentration: 5µg/ml.


### Species Reactivity

**Tested:** Human, Pig


**Expected from sequence similarity:** Human, Dog, Pig, Cow



EB12188 (0.1 $\mu$ g/ml) staining of K562 lysate (35 $\mu$ g protein in RIPA buffer). Detected by chemiluminescence.



EB12188 (0.3 $\mu$ g/ml) staining of Pig Testis lysate (35 $\mu$ g protein in RIPA buffer). Detected by chemiluminescence.



EB12188 (5µg/ml) staining of paraffin embedded Human Pancreas. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.