



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB06061 - Goat Anti-EMI2 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** FBXO43, EMI2, F-box protein 43, Fbx43, early mitotic inhibitor 2, ERP1

**Official Symbol:** FBXO43

**Accession Number(s):** NP\_001025031.2

**Human GeneID(s):** [286151](#)

**Non-Human GeneID(s):** 78803 (mouse), 315034 (rat)

### Immunogen

Peptide with sequence C-PGSAQSKRNLKRL, from the C Terminus of the protein sequence according to NP\_001025031.2.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Approx 80kDa band observed in Mouse Ovary lysates (calculated MW of 78.4kDa according to NP\_001025031.2). Recommended concentration: 0.3-1µg/ml.

Primary incubation was 1 hour. Preliminary testing was unsuccessful on Human, Rat and Pig Testis, and Human Ovary for this particular batch.

### Species Reactivity

**Tested:** Mouse

**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Pig, Cow

### Specific References

**This antibody (previous batch) has been successfully used in Western blot on Mouse:**

Pasternak M, Pfender S, Santhanam B, Schuh M

The BTG4 and CAF1 complex prevents the spontaneous activation of eggs by deadenylating maternal mRNAs.

Open Biol. 2016 Sep;6(9)

PMID: 27605379

**This antibody (previous batch) has been successfully used in Western blot on Mouse:**

Bernhardt ML, Kong BY, Kim AM, O'Halloran TV, Woodruff TK.


A zinc-dependent mechanism regulates meiotic progression in mammalian oocytes. Biol Reprod. 2012 Apr 19;86(4):114.

PMID: 22302686

**This antibody (previous batch) has been successfully used in Western blot, Immunocytochemistry and Flow Cytometry on Rat:**

Perrard MH, Chassaing E, Montillet G, Sabido O, Durand P.

Cytostatic Factor Proteins Are Present in Male Meiotic Cells and  $\beta$ -Nerve Growth Factor  
Increases Mos Levels in Rat Late Spermatocytes.  
PLoS One. 2009 Oct 5;4(10):e7237.  
PMID: 19802389



EB06061 (0.3 $\mu$ g/ml) staining of Mouse Ovary lysate (35 $\mu$ g protein in RIPA buffer). Detected by chemiluminescence.