



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB10691 - Goat Anti-Endophilin-A2 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** SH3GL1, SH3-domain GRB2-like 1, CNSA1, EEN, MGC111371, SH3D2B, SH3P8, EEN fusion partner of MLL, SH3 domain GRB2-like 1, SH3 domain-containing GRB2-like protein 1, SH3-containing Grb-2-like 1 protein, SH3-containing protein EEN, endophilin-A2, endophilin-2, SH3 domain protein 2B, extra 11-19 leukemia fusion, extra eleven-nineteen leukemia fusion gene protein

**Official Symbol:** SH3GL1

**Accession Number(s):** NP\_003016.1

**Human GeneID(s):** [6455](#)

### Immunogen

Peptide with sequence C-EPFDLGEPEQSN, from the internal region of the protein sequence according to NP\_003016.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.


**Western blot:** Approx 48kDa band observed in lysates of cell line K562 (calculated MW of 41.5kDa according to NP\_003016.1. The observed molecular weight corresponds to earlier findings in literature with different antibodies (So et al, Proc Natl Acad Sci U S A. 1997 Mar 18;94(6):2563-8.; PMID: 9122235). Recommended concentration: 0.3-1µg/ml. Primary incubation was 1 hour.

**IHC:** Paraffin embedded Human Small Intestine. Recommended concentration: 5µg/ml.


### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Dog, Cow



EB10691 (0.3µg/ml) staining of K562 lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB10691 (5µg/ml) staining of paraffin embedded Human Small Intestine. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.