



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB07144 - Goat Anti-EP1 receptor / PTGER1 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** PTGER1, prostaglandin E receptor 1 (subtype EP1), 42kDa, EP1, PGE receptor, EP1 subtype, prostaglandin E receptor 1, subtype EP1, prostanoid EP1 receptor

**Official Symbol:** PTGER1

**Accession Number(s):** NP\_000946.2

**Human GeneID(s):** [5731](#)

### Immunogen

Peptide with sequence C-DSRRRWGAHGPRS, from the internal region of the protein sequence according to NP\_000946.2.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested


**Peptide ELISA:** antibody detection limit dilution 1:128000.

**IHC:** In paraffin embedded Human Kidney shows membranous staining in the epithelial cells of DCT. Recommended concentration: 1-2µg/ml. Paraffin embedded Human Spleen. Recommended concentration: 5µg/ml.


### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human



EB07144 (2µg/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer pH 6, HRP-staining.



EB07144 (5µg/ml) staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.