

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

#### www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB07445 - Goat Anti-EPHB2 / DRT / Nuk Antibody

Size: 100µg specific antibody in 200µl



#### **Target Protein**

**Principal Names:** EPHB2, DRT, EPH receptor B2, Cek5, Qek5, Sek3, ETECK, Prkm5, EPHT3, ERK, Hek5, MGC87492, Tyro5, developmentally-regulated eph-related tyrosine kinase, elk-related tyrosine kinase, eph tyrosine kinase 3, ephrin receptor EphB2

Official Symbol: EPHB2

Accession Number(s): NP\_004433.2; NP\_059145.2

Human GeneID(s): 2048

Non-Human GenelD(s): 13844 (mouse)

Important Comments: This antibody is expected to recognise both reported isoforms (

NP\_059145.2 and NP\_004433.2).

## **Immunogen**

Peptide with sequence C-QKDRNHRPKFGQ, from the internal region of the protein sequence according to NP\_004433.2; NP\_059145.2.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:4000.

**Western blot:** Approx 60kDa band observed in Human Lung and Placenta lysates (calculated MW of 110kDa according to NP\_059145.2 and NP\_004433.2). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Litterst et al, J Biol Chem. 2007 Apr 10; PMID: 17428795). Recommended concentration: 1-3µg/ml.

## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog



EB07445 (1μg/ml) staining of Human Lung lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour.

Detected by chemiluminescence.