

#### International Office

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB09492 - Goat Anti-EPM2AIP1 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

Principal Names: EPM2AIP1, EPM2A (laforin) interacting protein 1, FLJ11207,

KIAA0766, EPM2A interacting protein 1

Official Symbol: EPM2AIP1

Accession Number(s): NP\_055620.1

Human GeneID(s): 9852

### **Immunogen**

Peptide with sequence C-TRNQHTLSQPLTDEH, from the internal region (near C Terminus) of the protein sequence according to NP\_055620.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:8000.

**Western blot:** In transfected HEK293 transiently expressing full-length Human EPM2AIP1 (myc and DYKDDDDK tagged), a band of approx. 70kDa was observed. No bands were observed in mock-transfected HEK293 and the same band was observed using anti-DYKDDDDK tag antibody. Recommended concentration, 1-2µg/ml.


**IHC:** In paraffin embedded Human Heart shows regular staining of sarcoplasmic reticulum in the cardiac fibers. Recommended concentration, 5-10µg/ml.

Purchase the +ve control! The transfected lysate used to QC this product is available for sale. Please email sales@everestbiotech.com to request datasheet, pricing and delivery information.


## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog



EB09492 (5μg/ml) staining of paraffin embedded Human Heart. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



HEK293 lysate (10ug protein in RIPA buffer) overexpressing Human EPM2AIP1 with DYKDDDDK tag probed with EB09492 (1ug/ml) in Lane A and probed with anti-DYKDDDK Tag (1/1000) in lane C. Mock-transfected HEK293 probed with EB09492 (1mg/ml) in Lane B. Primary incubations were for 1 hour. Detected by chemiluminescence.