

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB08306 - Goat Anti-EXO1 Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

Principal Names: EXO1, exonuclease 1, HEX1, hExol, rad2 nuclease family member,

homolog of S. cerevisiae exonuclease 1

Official Symbol: EXO1

Accession Number(s): NP\_003677.3; NP\_006018.3; NP\_569082.1

Human GeneID(s): 9156

Important Comments: This antibody is expected to recognise all three reported isoforms

(NP\_003677.3; NP\_006018.3 and NP\_569082.1)

### **Immunogen**

Peptide with sequence C-HRNYSPRPESGT, from the internal region of the protein sequence according to NP\_003677.3; NP\_006018.3; NP\_569082.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

#### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:64000.

**Western blot:** Approx 90kDa band observed in lysates in cell lines HEK293 and A549 (calculated MW of 89.1kDa according to NP\_003677.3). Recommended concentration: 0.2-0.5µg/ml. An additional band of unknown identity was also consistently observed at 38kDa. This band was successfully blocked by incubation with the immunising peptide. We would appreciate any feedback from people in the field - have any such results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?

# **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB08306 (0.2 $\mu$ g/ml) staining of A549 cell lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.