



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB09731 - Goat Anti-EYA1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: EYA1, eyes absent homolog 1 (Drosophila), BOP, BOR, MGC141875, Eyes absent, Drosophila, homolog of, 1, Melnick-Fraser syndrome, OTTHUMP00000195053, eyes absent 1

Official Symbol: EYA1

Accession Number(s): NP_742057.1; NP_000494.2; NP_742056.1

Human GeneID(s): [2138](#)

Non-Human GeneID(s): 14048 (mouse), 502935 (rat)

Important Comments: This antibody is expected to recognize all isoforms (NP_742057.1; NP_000494.2; NP_742056.1). Reported variants represent identical protein: NP_000494.2, NP_742055.1

Immunogen

Peptide with sequence C-TDPTAEYSTIHSP, from the internal region of the protein sequence according to NP_742057.1; NP_000494.2; NP_742056.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 55kDa band observed in lysates of cell line HEK293 (calculated MW of 60.7kDa according to NP_742056.1). Recommended concentration: 1-3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow



EB09731 (2 μ g/ml) staining of HEK293 lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.