

UK Office

Everest Biotech Ltd Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries: info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106 Ramona, CA 92065 USA

Inquiries: info@everestbiotech.com Sales: usasales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: 888-320-4628 (toll-free) Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB08049 - Goat Anti-Factor XIIIa (703-717) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: factor XIIIa, F13A1, coagulation factor XIII, A1 polypeptide, RP11-232H4.1, F13A, Coagulation factor XIII A chain, Coagulation factor XIII, A polypeptide, TGase, bA525O21.1 (coagulation factor XIII, A1 polypeptide), coagulation factor XIII A1 subunit Official Symbol: F13A1 Accession Number(s): NP_000120.2 Human GeneID(s): <u>2162</u>

Immunogen

Peptide with sequence C-HRKLIASMSSDSLRH, from the internal region of the protein sequence according to NP_000120.2.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000. **Western blot:** Approx 90kDa band observed in Human Placenta lysates (calculated MW of 83.3kDa according to NP_000120.2). Recommended concentration: 0.05-0.2µg/ml.

Species Reactivity

Tested: Human Expected from sequence similarity: Human, Mouse

| - | 250kDa 150kDa 100kDa |
|---|----------------------------|
| | 75kDa |
| | 50kDa |
| | 37kDa |
| | 25kDa |
| | 20kDa |
| | |

15kDa

EB08049 (0.1µg/ml) staining of Human Placenta lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.