

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# **EB10143 - Goat Anti-FANCE Antibody**

Size: 100µg specific antibody in 200µl



# **Target Protein**

Principal Names: FACE, FAE, Fanconi anemia, complementation group E, FANCE

Official Symbol: FANCE

Accession Number(s): NP\_068741.1

Human GeneID(s): 2178

### **Immunogen**

Peptide with sequence C-EHKSLESLADGGS, from the internal region of the protein sequence according to NP\_068741.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.


Western blot: Preliminary experiments gave an approx 70kDa band in Human Adipose and Uterus lysates after 0.1μg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 58.7kDa according to NP\_068741.1. The 70kDa band was successfully blocked by incubation with the immunizing peptide. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?

**IHC:** In paraffin embedded Human Placenta shows nuclear staining in some trophoblasts. Recommended concentration, 3-6µg/ml.

## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Cow



EB10143 (3.8 $\mu$ g/ml) staining of paraffin embedded Human Placenta. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.