

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB09626 - Goat Anti-FAS / CD95 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** FAS, Fas (TNF receptor superfamily, member 6), ALPS1A, APO-1, APT1, CD95, FAS1, FASTM, TNFRSF6, APO-1 cell surface antigen, CD95 antigen, Fas AMA, Fas antigen, OTTHUMP00000020045, OTTHUMP00000020046, OTTHUMP00000020051, OTTHUMP00000059646, apoptosis antigen 1, tumor necrosis factor receptor superfamily member 6, tumor necrosis factor receptor superfamily, member 6

**Official Symbol:** FAS

**Accession Number(s):** NP\_000034.1; NP\_690610.1; NP\_690611.1

**Human GeneID(s):** [355](#)

**Important Comments:** This antibody is expected to recognize reported isoforms 1 (NP\_000034.1), 2 (NP\_690610.1) and 3 (NP\_690611.1).

### Immunogen

Peptide with sequence KTCRKHRKENQGSH, from the internal region of the protein sequence according to NP\_000034.1; NP\_690610.1; NP\_690611.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Approx 37kDa band observed in lysates of cell line MOLT4 (calculated MW of 37.7kDa according to NP\_000034.1). Recommended concentration: 0.1-0.3µg/ml. Primary incubation was 1 hour.

**IHC:** Paraffin embedded Human Tonsil and Small Intestine. Recommended concentration: 3.75µg/ml.


### Species Reactivity

**Tested:** Human


**Expected from sequence similarity:** Human



EB09626 (0.1 $\mu$ g/ml) staining of MOLT4 lysate (35 $\mu$ g protein in RIPA buffer). Detected by chemiluminescence.



EB09626 ( $\mu$ g/ml) staining of paraffin embedded Human Tonsil. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB09626 (3.75µg/ml) staining of paraffin embedded Human Small Intestine. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.