

### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

## www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB12446 - Goat Anti-FCRL1 (aa165-177) Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** FCRL1, Fc receptor-like 1, CD307a, FCRH1, IFGP1, IRTA5, Fc receptor-like protein 1, IFGP family protein 1, fc receptor homolog 1, fcR-like protein 1, hIFGP1, immune receptor translocation-associated protein 5, immunoglobulin superfamily

Fc receptor, gp42

Official Symbol: FCRL1

Accession Number(s): NP\_443170.1; NP\_001152869.1; NP\_001152870.1

Human GenelD(s): 115350

Important Comments: This antibody is expected to recognize all reported isoforms

(NP\_443170.1; NP\_001152869.1; NP\_001152870.1).

# **Immunogen**

Peptide with sequence C-TAEYEIPSVRESD, from the internal region of the protein sequence according to NP\_443170.1; NP\_001152869.1; NP\_001152870.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**


Peptide ELISA: antibody detection limit dilution 1:64000.

**Western blot:** Approx 45 + 40kDa bands observed in Human Heart lysates and 38kDa in Human Liver lysates (calculated MW of 46.9kDa according to NP\_443170.1 and of 40.4kDa according to NP\_001152869.1). Recommended concentration: 0.3-1µg/ml.

### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human



EB12446 (0.3μg/ml) staining of Human Heart (A) and Liver (B) lysates (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.