

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

 $\underline{usasales@everest biotech.com}$

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB11836 - Goat Anti-FGF13 (isoform 1) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: FGF13, FGF-13, FGF2, FHF2, FHF-2, fibroblast growth factor 13,

fibroblast growth factor homologous factor 2

Official Symbol: FGF13

Accession Number(s): NP_004105.1

Human GenelD(s): 2258

Non-Human GenelD(s): 14168 (mouse)

Important Comments: This antibody is expected to recognize isoform 1 (NP_004105.1) only. Reported variants represent identical protein: NP_001132974.1, NP_001132973.1

Immunogen

Peptide with sequence KGKTSCDKNKLNVFS, from the internal region of the protein sequence according to NP_004105.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 28kDa band observed in Mouse Brain lysates (calculated MW of 27.6kDa according to NP_034330.2). Recommended concentration: 0.3-1µg/ml.

Species Reactivity

Tested: Mouse

Expected from sequence similarity: Human, Mouse, Dog, Pig, Cow



EB11836 ($0.5\mu g/ml$) staining of Mouse Brain lysate ($35\mu g$ protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.