



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB11834 - Goat Anti-Fgf14 (mouse N terminus) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: Fgf14, FGF-14, Fhf4, FHF-4, fibroblast growth factor 14, fibroblast growth factor homologous factor 4, mFHF-4(1B)

Official Symbol: Fgf14

Accession Number(s): NP_034331.2; NP_997550.1

Human GeneID(s): [2259](#)

Non-Human GeneID(s): 14169 (mouse), 63851 (rat)

Important Comments: This antibody is expected to recognize mouse isoform a (NP_034331.2) and human isoform 1A (NP_004106.1).

Immunogen

Peptide with sequence REQHWDRPSASR-C, from the N Terminus of the protein sequence according to NP_034331.2; NP_997550.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.


Western blot: Approx 28kDa band observed in Mouse and Human Brain lysates (calculated MW of 28.3kDa according to NP_997550.1 and 27.7kDa according to NP_034331.2). Recommended concentration: 1-3µg/ml. Primary incubation was 1 hour.

IHC: Cryosection of immersion-fixed (1% PFA/0.5% methanol) Human Hypothalamus. Optimal dilution: 1:5000. *Vibratome prepared sample for optimal labelling.


Species Reactivity

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow



EB11834 (1 μ g/ml) staining of Mouse Brain lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



EB11834 (1:5000) immunostaining of FGF14-positive axon initial segments in cryosection of an immersion-fixed (1% PFA/0.5% methanol) human hypothalamus*. Antigen retrieval with citrate buffer pH 6 at 80C for 30min. HRP-staining with Ni-DAB, after Biotin-SP-anti-goat (IgG) method. Data obtained by Drs. Éva Rumpler and Erik Hrabovszky, Inst, Exp, Med, Budapest, Hungary.