

International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB08456 - Goat Anti-FGFR2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: FGFR2, fibroblast growth factor receptor 2 (bacteria-expressed kinase, keratinocyte growth factor receptor, craniofacial dysostosis 1, Crouzon syndrome, Pfeiffer syndrome, Jackson-Weiss syndrome), BEK, BFR-1, CD332, CEK3, CFD1, ECT1, JWS, K-SAM

Official Symbol: FGFR2

Accession Number(s): NP_000132.3; NP_075259.4; NP_001138385.1; NP_001138386.1; NP_001138387.1; NP_001138388.1; NP_001138390.1; NP_001138391.1

Human GeneID(s): [2263](#)

Non-Human GeneID(s): 14183 (mouse), 25022 (rat)

Important Comments: This antibody is expected to recognise all reported isoforms except isoform 7 (NP_001138389.1). The immunizing peptide represents part of the extracellular domain.

Immunogen

Peptide with sequence C-GREKEITASPDY, from the internal region of the protein sequence according to NP_000132.3; NP_075259.4; NP_001138385.1; NP_001138386.1; NP_001138387.1; NP_001138388.1; NP_001138390.1; NP_001138391.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 100kDa band observed in Human lysates of cell line A549 (calculated MW of 92.0kDa according to NP_000132.1 and of 92.1kDa according to NP_075259.2). Recommended concentration: 0.3-1µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow



EB08456 (0.3 μ g/ml) staining of A549 lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.