



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06507 - Goat Anti-Fhl1/ Slim1 (mouse) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: FHL1, SLIM1, KYO-T, RP11-535K18.1, FHL1A, four and a half LIM domains 1, Four-and-a-half LIM domains 1, skeletal muscle LIM-protein1, FHL1B, KYOT, MGC111107, SLIMMER, XMPMA, four-and-a-half Lin11, Isl-1 and Mec-3 domains 1, FHL-1, FLH1A, SLIM-1, LIM protein SLIMMER, SLIM

Official Symbol: Fhl1

Accession Number(s): NP_001070830.1; NP_001274729.1

Non-Human GeneID(s): 14199 (mouse), 25177 (rat)

Important Comments: This antibody is expected to recognize Mouse NP_001070830.1 iso 2, NP_001274729.1 iso 3 and Rat NP_001029098.1 iso 1, NP_001258128.1 iso 3.

Immunogen

Peptide with sequence CNKRQVFHNEQVY, from the internal region of the protein sequence according to NP_001070830.1; NP_001274729.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 32kDa band observed in Human, Mouse and Rat Skeletal Muscle lysates and in Rat Lung and Testis lysates (calculated MW of 31.9kDa according to Mouse NP_001274729.1 and Rat NP_001258128.1). Recommended concentration: 0.01-0.5µg/ml. Primary incubation 1 hour at room temperature.


Immunofluorescence: Strong expression of the protein seen in the cytoplasm of U2OS cells. Recommended concentration: 10µg/ml.

Flow Cytometry: Flow cytometric analysis of U2OS cells. Recommended concentration: 10µg/ml.


Species Reactivity

Tested: Human, Mouse, Rat


Expected from sequence similarity: Human, Mouse, Rat, Ciow




EB06507 (0.1 μ g/ml) staining of Mouse Skeletal Muscle (A), Rat Lung (B), Rat Skeletal Muscle (C) and (0.5 μ g/ml Rat Testes (D) lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



EB06507 (0.01 μ g/ml) staining of Human Skeletal Muscle lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



EB06507 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB06507 Flow cytometric analysis of paraformaldehyde fixed U2OS cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml).