



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07434 - Goat Anti-FLAP / ALOX5AP (internal) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: ALOX5AP, FLAP, arachidonate 5-lipoxygenase-activating protein, MK-886-binding protein, five-lipoxygenase activating protein

Official Symbol: ALOX5AP

Accession Number(s): NP_001620.2; NP_001191335.1

Human GeneID(s): [241](#)

Important Comments: This antibody is expected to recognize both reported isoforms (NP_001620.2; NP_001191335.1).

Immunogen

Peptide with sequence C-HKVEHESRTQNGR, from the internal region of the protein sequence according to NP_001620.2; NP_001191335.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 16kDa band observed in lysates of U937 and of Peripheral Blood Mononucleocytes (calculated MW of 18.2kDa according to NP_001620.2). Recommended concentration: 0.1-0.3µg/ml. Primary incubation was 1 hour.

IHC: In paraffin embedded Human Spleen shows staining of select splenocytes. Paraffin embedded Human Lung and Small Intestine. Recommended concentration: 3.75µg/ml.


Species Reactivity

Tested: Human


Expected from sequence similarity: Human, Dog




EB07434 (0.1 μ g/ml) staining of Human U937 cell lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



EB07434 (3.75 μ g/ml) staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB07434 (3.75µg/ml) staining of paraffin embedded Human Lung. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB07434 (3.75µg/ml) staining of paraffin embedded Human Small Intestine. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.