



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB07434-B - Goat Anti-FLAP / ALOX5AP (internal), Biotinylated Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** ALOX5AP, arachidonate 5-lipoxygenase-activating protein, FLAP, MK-886-binding protein, five-lipoxygenase activating protein

**Official Symbol:** ALOX5AP

**Accession Number(s):** NP\_001620.2; NP\_001191335.1

**Human GeneID(s):** [241](#)

**Important Comments:** This antibody is expected to recognize both reported isoforms (NP\_001620.2; NP\_001191335.1).

### Immunogen

Peptide with sequence C-HKVEHESRTQNGR., from the internal region of the protein sequence according to NP\_001620.2; NP\_001191335.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested


**Peptide ELISA:** antibody detection limit dilution 1:128000.

**Western blot:** Approx 16kDa band observed in lysates of cell line U937 (calculated MW of 18.2kDa according to NP\_001620.2). See non-biotinylated parental product's datasheet for further QC data. Recommended concentration: 0.1-0.3µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Dog



Biotinylated EB07434 (0.3 $\mu$ g/ml) staining of U937 lysate (35 $\mu$ g protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.