

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB06022 - Goat Anti-FOXL2 / BPES Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: FOXL2, BPES, forkhead box L2, PFRK, BPES1, PINTO, forkhead transcription factor FOXL2, blepharophimosis, epicanthus inversus and ptosis,

Blepharophimosis, epicanthus inversus, and ptosis 1, POF3

Official Symbol: FOXL2

Accession Number(s): NP_075555.1

Human GeneID(s): 668

Non-Human GenelD(s): 26927 (mouse)

Immunogen

Peptide with sequence C-DSKTGALHSRLDL, from the C Terminus of the protein sequence according to NP_075555.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx. 50kDa band observed in Human and Mouse Ovary lysates (calculated MW of 38.8kDa according to Human NP_075555.1 and Mouse NP_036150.1).

This molecular weight is observed by other Sources. Recommended concentration

1-3ug/ml. Primary incubation 1 hour at room temperature.

IHC: Paraffin embedded Human Ovary. Recommended concentration: 5µg/ml.

Species Reactivity

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat, Cow

Specific Reference


This antibody (previous batch) has been successfully used in IF on Mouse:

Padua MB, Fox SC, Jiang T, Morse DA, Tevosian SG.

Simultaneous Gene Deletion of Gata4 and Gata6 Leads to Early Disruption of Follicular Development and Germ Cell Loss in the Murine Ovary.


Biol Reprod. 2014 Jun 4.

PMID: 24899573




EB06022 (2μg/ml) staining of Human (A) and (1μg/ml) Mouse (B) Ovary lysate (35μg protein in RIPA buffer).

Detected by chemiluminescence.



EB06022 ($5\mu g/ml$) staining of paraffin embedded Human Ovary. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



 ${\sf EB06022\ Negative\ Control\ showing\ staining\ of\ paraffin\ embedded\ Human\ Ovary,\ with\ no\ primary\ antibody.}$