



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05226-T - Goat Anti-FOXP2 (C terminus) Antibody - Trial

Size: 20µg specific antibody in 40µl



Target Protein

Principal Names: FOXP2, forkhead box P2, SPCH1, CAGH44, TNRC10, CAG repeat protein 44, speech and language disorder 1, trinucleotide repeat containing 10, forkhead/winged-helix transcription factor, DKFZp686H1726, OTTHUMP00000196932

Official Symbol: FOXP2

Accession Number(s): NP_055306.1; NP_683696.2; NP_683697.1

Human GeneID(s): [93986](#)

Important Comments: This antibody is expected to recognise all three reported isoforms (NP_055306.1; NP_683696.2; NP_683697.1).

Immunogen

Peptide with sequence C-REIEEEPLSEDL, from the C Terminus of the protein sequence according to NP_055306.1; NP_683696.2; NP_683697.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 80-90kDa band observed in Human Brain (Cerebellum) lysates (calculated MW of 82.6kDa according to NP_683696.2). This molecular weight is routinely observed by other sources. Recommended concentration: 0.5-2µg/ml. Primary incubation 1 hour at room temperature.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow

Specific References

This antibody has been successfully used in the following paper:

Amber G. Cook, Taylor V. Bishop, Hannah R. Crowe, Daniel N. Stevens, Lauren Reine, Alexandra L. Joyner and Andrew K. Lawton

Cell division angle predicts the level of tissue mechanics that tune the amount of cerebellar folding.

Development. 2024 Feb 1;151(3):dev202184.

PMID: 38251865

This antibody (previous batch) has been successfully used in ICC on Mouse:

Yuanjun Luo, Yuhan Chao, Jingyun Zhang, Tatsumi Hirata and Izumi Sugihara
Neurogenic timing of the inferior olive subdivisions is related to the olivocerebellar projection topography

Research Square, August 2022, [<https://doi.org/10.21203/rs.3.rs-1923633/v1>]

PMID: 37130860

This antibody (previous batch) has been successfully used in ICC on Mouse:

Khoa Tran-Anh, Jingyun Zhang, Viet Tuan Nguyen-Minh, Hirofumi Fujita, Tatsumi Hirata and Izumi Sugihara

Common Origin of the Cerebellar Dual Somatotopic Areas Revealed by Tracking Embryonic Purkinje Cell Clusters with Birthdate Tagging

eNeuro. 2020 Dec 14;7(6):ENEURO.0251-20.2020.

PMID: 33055198

This antibody (previous batch) has been successfully used in IF on Mouse:

Martinez-Chavez E, Scheerer C, Wizenmann A, Blaess S

The zinc-finger transcription factor GLI3 is a regulator of precerebellar neuronal migration. Development. 2018 Dec 17;145(24). pii: dev166033.

PMID: 30470704

This antibody (previous batch) has been successfully used in ICC on Mouse:

Gideon A Sarpong, Suteera Vibulyaseck, Yuanjun Luo, Mohammad S Biswas, Hirofumi Fujita, Shinji Hirano, Izumi Sugihara

Cerebellar modules in the olivo-cortico-nuclear loop demarcated by pcdh10 expression in the adult mouse

J Comp Neurol. 2018 Oct 15;526(15):2406-2427

PMID: 30004589

This antibody (previous batch) has been successfully used in IF on Mouse:

Vibulyaseck S, Fujita H, Luo Y, Tran AK, Oh-Nishi A, Ono Y, Hirano S, Sugihara I

Spatial rearrangement of Purkinje cell subsets forms the transverse and longitudinal compartmentalization in the mouse embryonic cerebellum.

J Comp Neurol. 2017 Oct 1;525(14):2971-2990.

PMID: 28542916

This antibody (previous batch) has been successfully used in IHC on Human and Mouse:

Haldipur P, Dang D, Aldinger KA, Janson OK, Guimiot F, Adle-Biasette H, Dobyns WB, Siebert JR, Russo R, Millen KJ.

Phenotypic outcomes in Mouse and Human Foxc1 dependent Dandy-Walker cerebellar malformation suggest shared mechanisms.

Elife. 2017 Jan 16;6. pii: e20898. doi: 10.7554/eLife.20898.

PMID: 28092268

This antibody (previous batch) has been successfully used in WB and IHC on Chicken:

Vibulyaseck S, Luo Y, Fujita H, Oh-Nishi A, Ohki-Hamazaki H, Sugihara I

Compartmentalization of the chick cerebellar cortex based on the link between the striped expression pattern of aldolase C and the topographic olivocerebellar projection

J Comp Neurol. 2015 Sep 1;523(13):1886-912.

PMID: 25732420

This antibody (previous batch) has been successfully used in IHC on Mouse:

Fujita H, Morita N, Furuichi T, Sugihara I.

Clustered fine compartmentalization of the mouse embryonic cerebellar cortex and its rearrangement into the postnatal striped configuration.

J Neurosci. 2012 Nov 7;32(45):15688-703.

PMID: 23136409


This antibody (previous batch) has been successfully used in IHC on Mouse:

Fujita H, Sugihara I.

FoxP2 expression in the cerebellum and inferior olive: Development of the transverse stripe-shaped expression pattern in the mouse cerebellar cortex.

J Comp Neurol. 2012 Feb 15;520(3):656-77. doi: 10.1002/cne.22760.

PMID: 21935935



EB05226 (2µg/ml) staining of Human Cerebellum lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.