

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB08348 - Goat Anti-Frizzled 7 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

Principal Names: rizzled 7, frizzled homolog 7 (Drosophila), FZD7, FzE3, Frizzled,

drosophila, homolog of, 7 **Official Symbol:** FZD7

Accession Number(s): NP\_003498.1

Human GeneID(s): 8324

Non-Human GenelD(s): 14369 (mouse)

### **Immunogen**

Peptide with sequence C-RFYHRLSHSSKGET, from the C Terminus of the protein sequence according to NP\_003498.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:64000.

**Western blot:** Approx 60kDa band observed in lysates of cell lines HEK293 and HepG2 (calculated MW of 63.6kDa according to NP\_003498.1). Recommended concentration:

IHC: Paraffin embedded Human Spleen. Recommended concentration: 5µg/ml.

## **Species Reactivity**


Tested: Human

Expected from sequence similarity: Human, Mouse



EB08348 (0.3μg/ml) staining of HepG2 cell lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour.

Detected by chemiluminescence.



EB08348 (5µg/ml) staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.