

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB08348 - Goat Anti-Frizzled 7 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: rizzled 7, frizzled homolog 7 (Drosophila), FZD7, FzE3, Frizzled,

drosophila, homolog of, 7 **Official Symbol:** FZD7

Accession Number(s): NP_003498.1

Human GeneID(s): 8324

Non-Human GenelD(s): 14369 (mouse)

Immunogen

Peptide with sequence C-RFYHRLSHSSKGET, from the C Terminus of the protein sequence according to NP_003498.1.

Please note the <u>peptide</u> is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 60kDa band observed in lysates of cell lines HEK293 and HepG2 (calculated MW of 63.6kDa according to NP_003498.1). Recommended concentration:

IHC: Paraffin embedded Human Spleen. Recommended concentration: 5µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse



EB08348 ($0.3\mu g/ml$) staining of HepG2 cell lysate ($35\mu g$ protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB08348 (5µg/ml) staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.