



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07361-T - Goat Anti-FXR1 Antibody - Trial

Size: 20µg specific antibody in 40µl



Target Protein

Principal Names: FXR1, fragile X mental retardation, autosomal homolog 1, fragile X mental retardation-related protein 1

Official Symbol: FXR1

Accession Number(s): NP_005078.2; NP_001013456.1; NP_001013457.1

Human GeneID(s): [8087](#)

Non-Human GeneID(s): 14359 (mouse), 361927 (rat)

Important Comments: This antibody is expected to recognise all reported isoforms (NP_005078.2, NP_001013456.1 and NP_001013457.1).

Immunogen

Peptide with sequence C-RIEGDNENKLPRED, from the internal region of the protein sequence according to NP_005078.2; NP_001013456.1; NP_001013457.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 75+80kDa band observed in lysates of cell line NIH-3T3 (calculated MW of 59.9kDa according to NP_001013457.1). The observed molecular weight corresponds to earlier findings in literature (Khandjian et al Hum Mol Genet. 1998 Dec;7(13):2121-8.; PMID: 9817930). An additional band of unknown identity was also consistently observed at 37kDa. This band was successfully blocked by incubation with the immunizing peptide. Recommended concentration. 1-3µg/ml. Primary incubation 1 hour at room temperature.

Immunofluorescence: Strong expression of the protein seen in the cytoplasm of A431 and HeLa cells. Recommended concentration: 10µg/ml.

Flow Cytometry: Flow cytometric analysis of HeLa cells. Recommended concentration: 10ug/ml.


Species Reactivity

Tested: Human, Mouse


Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow




EB07361 (1 μ g/ml) staining of NIH-3T3 cell lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



EB07361 Immunofluorescence analysis of paraformaldehyde fixed A431 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB07361 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB07361 Flow cytometric analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.