

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB05153 - Goat Anti-FYN Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** FYN, SLK, SYN, MGC45350, FYN oncogene related to SRC, FGR, YES, src-like kinase, c-syn protooncogene, tyrosine kinase p59fyn(T), src/yes-related novel gene, OKT3-induced calcium influx regulator, proto-oncogene tyrosine-protein kinase fyn, RP1-66H14.1, OTTHUMP0000017917, protein-tyrosine kinase fyn, src-like kinase

**Official Symbol:** FYN

**Accession Number(s):** NP\_002028.1; NP\_694592.1; NP\_694593.1

**Human GeneID(s):** [2534](#)

**Non-Human GeneID(s):** 14360 (mouse), 25150 (rat)

**Important Comments:** This antibody is expected to recognise all three reported isoforms (as represented by NP\_002028.1; NP\_694592.1; NP\_694593.1).

### Immunogen

Peptide with sequence VQCKDKEATKLTE, from the N Terminus of the protein sequence according to NP\_002028.1; NP\_694592.1; NP\_694593.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested


**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Approx 55kDa band observed in lysates of cell lines A431 and Jurkat (calculated MW of 54.5kDa according to NP\_694593.1). Recommended for use at 0.3-1µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Pig, Cow



EB05153 (0.3 $\mu$ g/ml) staining of A431 (A) and Jurkat lysate (B) (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.