

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB07841 - Goat Anti-G6PD (aa 308 - 320) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: G6PD, glucose-6-phosphate dehydrogenase, G6PD1,

glucose-6-phosphate 1-dehydrogenase, glucose-6-phosphate dehydrogenase, G6PD

Official Symbol: G6PD

Accession Number(s): NP_000393.4; NP_001035810.1

Human GeneID(s): 2539

Non-Human GenelD(s): 14381 (mouse), 24377 (rat)

Important Comments: This antibody is expected to recognise both reported isoforms

(NP_000393.4 and NP_001035810.1).

Immunogen

Peptide with sequence C-STNSDDVRDEKVK, from the internal region of the protein sequence according to NP_000393.4; NP_001035810.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 55-60kDa double band observed in Human Testis, Thyroid and Tonsil lysates (calculated MW of 62.5kDa according to NP_000393.4 and 59.3kDa according to NP_001035810.1). Recommended concentration: 0.03-0.1µg/ml. Primary incubation was 1 hour.

IHC: Paraffin embedded Human Testis and Spleen. Recommended concentration: 2.5µg/ml.

Additional validation: This antibody has been successfully used in the following paper: Sikorski et al. (2018) PMID: 30377371.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog

Specific References

This antibody has been successfully used in the following paper:

Adriana Ann Garcia, Irimpan I. Mathews, Naoki Horikoshi, Tsutomu Matsui, Manat Kaur, Soichi Wakatsuki, and Daria Mochly-Rosen

Stabilization of glucose-6-phosphate dehydrogenase oligomers enhances catalytic activity and stability of clinical variants.

J Biol Chem. 2022 Mar; 298(3): 101610.

PMID: 35065072

This antibody has been successfully used in the following paper:


Krzysztof Sikorski, Adi Mehta, Marit Inngjerdingen, Flourina Thakor, Simon Kling, Tomas Kalina, Tuula A. Nyman, Maria Ekman Stensland, Wei Zhou, Gustavo A. De Souza, Lars

Holden, Jan Stuchly, Markus Templin and Fridtjof Lund-Johansen A high-throughput pipeline for validation of antibodies Nat Methods. 2018 Nov;15(11):909-912


PMID: 30377371

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB07841 ($0.03\mu g/ml$) staining of Human Testis lysate ($35\mu g$ protein in RIPA buffer). Detected by chemiluminescence.



EB07841 (2.5 μ g/ml) staining of paraffin embedded Human Testis. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB07841 (2.5μg/ml) staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.