

#### **UK Office**

### **Everest Biotech Ltd**

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

**Enquiries:** 

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

### **US Office**

### **Everest Biotech c/o Abcore**

405 Maple Street, Suite A106 Ramona,

CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

 $\underline{usasales@everest biotech.com}$ 

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB07710 - Goat Anti-Galectin 3 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** LGALS3, lectin, galactoside-binding, soluble, 3, CBP35, GAL3, GALBP, GALIG, LGALS2, MAC2, IgE-binding protein, MAC-2 antigen,

carbohydrate-binding protein 35, galactose-specific lectin 3, galectin 3, galectin-3 internal gene, laminin-binding protein, lectin, galactoside-binding, soluble, 3 (galectin 3)

Official Symbol: LGALS3

Accession Number(s): NP\_002297.2

Human GeneID(s): 3958

### **Immunogen**

Peptide with sequence CNTKLDNNWGREERQ, from the internal region of the protein sequence according to NP\_002297.2.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:128000.


**Western blot:** Approx 30kDa band observed in Human and Mouse Colon lysates (calculated MW of 26.2kDa according to NP\_002297.2). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Lotz et al, Proc Natl Acad Sci U S A. 1993 Apr 15;90(8):3466-70. PMID: 7682704). Recommended

concentration: 0.3-1µg/ml.

### **Species Reactivity**

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat



EB07710 (0.3 $\mu$ g/ml) staining of Human (A) and Mouse (B) Colon lysates (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.