



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB05571 - Goat Anti-GATA1 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** GATA1, GATA binding protein 1 (globin transcription factor 1), GF1, NFE1, ERYF1

**Official Symbol:** GATA1

**Accession Number(s):** NP\_002040.1

**Human GeneID(s):** [2623](#)

### Immunogen

Peptide with sequence C-PTTSTTVVAPLSS, from the C Terminus of the protein sequence according to NP\_002040.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Approx 50kDa band observed in nuclear lysates of cell line K562 (calculated MW of 42.8kDa according to NP\_002040.1). The observed molecular weight corresponds to different antibodies from other sources and was successfully blocked by incubation with the immunizing peptide. Recommended concentration: 1-3µg/ml. Primary incubation 1 hour at room temperature. **Negative Control:** Human Hippocampus lysate.


**Immunofluorescence:** Strong expression of the protein seen in the nuclei of HeLa cells. Recommended concentration: 10µg/ml.

**Flow Cytometry:** Flow cytometric analysis of K562 cells. Recommended concentration: 10ug/ml.


### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human




EB05571 (2 $\mu$ g/ml) staining of K562 nuclear cell lysate (A) + peptide (B) and negative control Human Hippocampus (C) lysate (35 $\mu$ g protein in RIPA buffer). Detected by chemiluminescence.



EB05571 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 $\mu$ g/ml) followed by Alexa Fluor 488 secondary antibody (2 $\mu$ g/ml), showing nuclear staining. Actin filaments were stained with phalloidin (red) and the nuclear stain is DAPI (blue). Negative control:

Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB05571 Flow cytometric analysis of paraformaldehyde fixed K562 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.