

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB06874 - Goat Anti-GATA1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: GATA1, GATA binding protein 1 (globin transcription factor 1), HGNC:4170, ERYF1, GF1, NFE1, GATA binding protein 1, GATA-binding protein 1 (globin transcription factor 1), erythrold transcription factor 1, globin transcription factor 1, transcription factor GATA1, GF-1, XLTT, NF-E1 DNA-binding protein, erythroid

transcription factor 1
Official Symbol: GATA1

Accession Number(s): NP_002040.1

Human GenelD(s): 2623

Immunogen

Peptide with sequence C-DAEAYRHSPVFQ, from the internal region of the protein sequence according to NP_002040.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 50-55kDa band observed in nuclear lysates of cell line K562 calculated MW of 42.8kDa according to NP_002040.1). This molecular weight is routinely observed by other sources. Recommended concentration: 0.3-1µg/ml. Primary incubation 1 hour at room temperature. **Negative Control:** Human Hippocampus lysate.


Immunofluorescence: Strong expression of the protein seen in the nuclei of HeLa and NIH3T3 cells. Recommended concentration: 10µg/ml.

Flow Cytometry: Flow cytometric analysis of K562 cells. Recommended concentration: 10ug/ml.


Species Reactivity

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow




EB06874 (1μg/ml) staining of K562 nuclear cell lysate (A) and negative control Human Hippocampus (B) lysate (35μg protein in RIPA buffer). Detected by chemiluminescence.




EB06874 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear staining. Actin filaments were stained with phalloidin (red) and the nuclear stain is DAPI (blue). Negative control:

Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB06874 Immunofluorescence analysis of paraformaldehyde fixed NIH3T3 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB06874 Flow cytometric analysis of paraformaldehyde fixed K562 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.